



Information

Tolerance of Field Grown Conifers to Herbicides

Bold text indicates information derived from companies' own literature.

Plain text indicated information derived from other sources.

Only a limited number of species and cultivars have been tested within each genus, it is therefore, essential to carry out small scale trials on your own plants prior to full scale treatment.

	Axit GR ¹	Butisan S	Dacthal w-75 ²	Devrinol	Diuron ³	Flexidor 125 ⁴	Kerb 50W ⁵	Ronstar liquid	Simazine ⁶	Venzar ⁷
Abies			T	T	T	T	T	T	T	
Araucaria						T			T	
Cedrus					T	T	T	T	M	
Chamaecyparis		T	T	T	T	T	T	T	M	
Cryptomeria						T		T	T	
Cunninghamia									T	
x Cupressocyparis				T		T	T	T	T	
Cupressus			T	T		T	T	T	T	
Ginkgo					T				T	
Juniperis		T	T	T		T	T	T	T	T
Larix					T	T	T		M	
Libocedrus									T	
Metasequoia									M	
Phyllocladus									T	
Picea		T	T	T	T	T	T	T	M	T
Pinus			S	T	T	T	T	T	T	T
Pseudotsuga	T				T	T	T		T	
Sequoia									T	
Sequoiadendron									T	
Taxus			T	T	T	T	T	T	M	
Thuja			T	T		T	T	T	M	T
Tsuga						T	T		T	

Key to table

S = Susceptible (unacceptable damage normally occurs – do not treat)

M = Moderately Susceptible (some damage may occur, but plants normally grow anyway)

T = Tolerant (damage does not normally occur)