



## Information

### Tolerance of Field Grown Conifers to Herbicides

Bold text indicates information derived from companies' own literature.  
Plain text indicated information derived from other sources.

Only a limited number of species and cultivars have been tested within each genus, it is therefore, essential to carry out small scale trials on your own plants prior to full scale treatment.

#### Key to table

- S = Susceptible (unacceptable damage normally occurs – do not treat) \* EAMU required  
M = Moderately Susceptible (some damage may occur, but plants normally grow anyway)  
T = Tolerant (damage does not normally occur)

	Butisan S	Dual Gold*	Flexidor	Kerb Flo	pendimethalin	Sumimax*	Venzar 500 SC
Abies		T	T	T	T	T	
Araucaria			T				
Cedrus			T	T			
Chamaecyparis	T		T	T	T		
Cryptomeria			T		T		
Cunninghamia							
x Cupressocyparis			T	T	T		
Cupressus			T	T	T		
Ginkgo		T			T		
Juniperus	T	T	T	T	T	T	T
Larix	T		T	T			
Libocedrus							
Metasequoia					T		
Phyllocladus							
Picea	T	T	T	T	T	T	T
Pinus	M	T	T	T	T	T	T
Pseudotsuga	T		T	T	T		
Sequoia					T		
Sequoiadendron					T		
Taxodium		T	T		T		
Taxus		T	T	T	T	T	
Thuja	T	T	T	T	T	T	T
Tsuga		T	T	T	T	T	

**Use plant protection products safely. Always read the label and product information before use**

Dove Associates shall in no event be liable for any loss or damage caused by the use of products mentioned in this document