



**DOVE**  
**ASSOCIATES**

**Horticultural Consultants**

Weggs Farm

Common Road

Dickleburgh, DISS

Norfolk IP21 4PJ

Tel: 01379 741200

Fax: 01379 741800

Email: [info@dovebugs.co.uk](mailto:info@dovebugs.co.uk)

[www.dovebugs.co.uk](http://www.dovebugs.co.uk)

## Information

### Scab Risk Periods

Close monitoring of weather conditions in the orchard and reference to Mills Period or Smith Period tables will provide a guide as to when periods of scab infection are either imminent or have already occurred and hence increased spray frequencies are required.

#### Mills Period Table

Mean Temperature		Hours of wet conditions required for scab infection of various severity to develop		
°C	(°F)	Light Infection	Moderate Infection	Severe Infection
6	(43)	30	40	60
8	(46)	19	25	38
10	(50)	14	19	29
12	(54)	11	16	24
14	(57)	10	14	21
16	(61)	9	12	19
18	(64)	9	12	18

**Example:** If wet conditions prevail, at a mean temperature of 12°C (54°F) for 24 hours, then severe scab infection is probable.

#### Smith Period Table

Apple scab infections are likely to occur if any one of the following conditions apply:

Rain -----	Combined with temperature		
	At 0800hrs	or	1700hrs
1. Starts to fall during the day and still falling at 1700hrs			10°C (50°F) or above
2. Starts to fall during the evening or night and leaves still wet at 0800hrs	10°C (50°F) or above		
3. Falls continuously for 12 hours or more	8°C (46°F) or above	or	8°C (46°F) or above
4. Falls continuously for 24 hours or more	7°C (44°F) or above	or	7°C (44°F) or above
5. Falls continuously for 36 hours or more	2°C (36°F) or above	or	2°C (36°F) or above

**Read the label before you buy – use pesticides safely**

Dove Associates shall in no event be liable for any loss or damage caused by the use of products mentioned in this document

© Dove Associates 05/02/09

The following products have either curative or protective qualities in respect of scab. For crops grown under protection check that the product you have selected is approved for application on protected crops. Some EC products can cause foliar damage if used in hot or dry conditions. Spray programmes for scab control should include an eradicant and protectant product. Some products can be tank mixed but always check the label. Application rates should be adjusted according to the frequency of application.

<b>Control product</b>	<b>[F]ield [G]lass [E]radicant [P]rotectant</b>	<b>FRAC code</b>
Amistar* <sup>+</sup>	F/G     E/P	11
Cercobin WG* <sup>+</sup>	F/G     E/P	1
Dithianon Flowable	F         E/P	M9
Flamenco	F         E/P	3
Frupica SC*	F         P	9
Indar 5EW	F         E/P	3
Plover <sup>+</sup>	F         E/P	3
Radspor FL	F         E/P	M7
Rubigan	F/G     E/P	3
Scala*	F/G     P	9
Stroby WG*	F/G     E/P	11
Sulphur Flowable	F/G     E	M2
Swift SC <sup>+</sup>	F         E/P	11
Sythane 20 EW	F/G     P	3
Topas	F         P	3
Triptam	F/G     P	1
Twist	F         E/P	11

\* LERAP applies to this product.

Avoid using products as successive treatments with the same FRAC code e.g. Amistar, Stroby WG, Swift SC and Twist.

The Long-Term Arrangements for Extension of Use have now ended. Dithianon Flowable, Flamenco, Frupica SC, Indar 5EW, Radspor FL, Scala, Sulphur Flowable and Twist will no longer be approved for use in ornamental plant production after 1 June 2009.

<sup>+</sup> = Amistar and Swift SC will require a new Specific Off-Label Approval from 1 June 2009 onwards for use in ornamental plant production. Go to [www.pesticides.gov.uk](http://www.pesticides.gov.uk) to download the information.

### **Plant subjects susceptible to scab**

Malus  
Pyrus  
Sorbus  
Crataegus  
Pyracantha  
Chaenomeles

### **Read the label before you buy – use pesticides safely**

Dove Associates shall in no event be liable for any loss or damage caused by the use of products mentioned in this document

© Dove Associates 05/02/09